

1. Tsuboi H, Segawa S, Yagishita M, Toko H, Honda F, Kitada A, Miki H, Ohyama A, Hagiwara S, Kondo Y, Matsumoto I, Sumida T.:
Activation mechanisms of monocytes/macrophages in adult-onset Still disease.
Front Immunol. (in press)
2. Furuyama K, Kondo Y, Kaneko S, Shimizu M, Tanimura R, Tsuboi H, Sumida T, Matsumoto I.:
Effector Th17-like regulatory T cells associate with the pathogenesis of rheumatoid arthritis.
Rheumatology (Oxford). (in press)
3. Konishi K, Nakamura K, Hanada Y, Kitanaga Y, Kubo S, Kinugasa F, Yamajuku D, Maeda M, Yamamoto N, Minematsu T, Ohbuchi M, Kondo Y, Sumida T.:
Preclinical Characterization of ASP2713, a Novel Ig β and Fc γ RIIB Cross-linking Antibody, for Prediction of Human Pharmacokinetics and Clinically Effective Dose.
J Pharm Sci. (Online ahead of print)
4. Kimura N, Kawahara T, Uemura Y, Atsumi T, Sumida T, Mimura T, Kawaguchi Y, Amano H, Iwasaki Y, Kaneko Y, Matsui T, Muro Y, Imura Y, Kanda T, Tanaka Y, Kawakami A, Jinnin M, Ishii T, Hiromura K, Miwa Y, Nakajima H, Kuwana M, Nishioka Y, Morinobu A, Kameda H, Kohsaka H.:
Branched chain amino acids in the treatment of polymyositis and dermatomyositis: a phase II/III, multi-center, randomized controlled trial.
Rheumatology (Oxford). (Online ahead of print)
5. Ichimura Y, Konishi R, Shobo M, Inoue S, Okune M, Maeda A, Tanaka R, Kubota N, Matsumoto I, Ishii A, Tamaoka A, Shimbo A, Mori M, Morio T, Kishi T, Miyamae T, Tanboon J, Inoue M, Nishino I, Fujimoto M, Nomura T, Okiyama N.:
Reliability of antinuclear matrix protein 2 antibody assays in idiopathic inflammatory myopathies is dependent on target protein properties.
J Dermatol. 49(4):441-447,2022
6. Furuyama K, Kondo Y, Shimizu M, Yokosawa M, Segawa S, Iizuka A, Tanimura R, Tsuboi H, Matsumoto I, Sumida T.:
ROR γ t +Foxp3 + regulatory T cells in the regulation of autoimmune arthritis.
Clin Exp Immunol. 207(2):176-187,2022
7. Ichimura Y, Konishi R, Shobo M, Inoue S, Okune M, Maeda A, Tanaka R, Kubota N, Matsumoto I, Ishii A, Tamaoka A, Shimbo A, Mori M, Morio T, Kishi T, Miyamae T, Tanboon J, Inoue M, Nishino I, Fujimoto M, Nomura T, Okiyama N.:
Anti-nuclear matrix protein 2 antibody-positive inflammatory myopathies represent extensive myositis without dermatomyositis-specific rash.
Rheumatology (Oxford). 61(3):1222-1227,2022